

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 January 2004 (15.01.2004)

PCT

(10) International Publication Number
WO 2004/006450 A1

(51) International Patent Classification⁷: **H04B 1/10, 1/69**

(21) International Application Number:
PCT/FI2003/000536

(22) International Filing Date: **2 July 2003 (02.07.2003)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
20021317 4 July 2002 (04.07.2002) FI

(71) Applicant (for all designated States except US): **NOKIA CORPORATION [FI/FI]; Keilalahdentie 4, FIN-02150 Espoo (FI).**

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SAARNISAARI,**

Harri [FI/FI]; Pihkatie 4 B 13, FIN-90800 Oulu (FI). HENTTU, Pertti [FI/FI]; Kaivosojantie 83, FIN-90470 Varjakka (FI).


(74) Agent: **KOLSTER OY AB; Iso Roobertinkatu 23, P.O. Box 148, FIN-00121 Helsinki (FI).**

(81) Designated States (*national*): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK (utility model), SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO utility model (GH), ARIPO patent (GH), ARIPO utility model (GM), ARIPO

[Continued on next page]

(54) Title: **SIGNAL-PROCESSING METHOD AND A RECEIVER**



(57) Abstract: The invention relates to a method for processing a signal, in which method a sample set is formed (300) and a set of absolute values is formed (302). The method comprises A) arranging (304) the elements of the absolute value set in the ascending order; B) setting (306) a threshold value; C) determining (308) the number of elements of a reference set to be formed; D) forming (309) a reference set; E) computing (310) a reference value by multiplying the mean or median of the reference set by the threshold value; F) comparing (312) the greatest element of the reference set with the reference value; G) increasing (316) the number of reference set elements; H) reiterating the preceding steps D to G until the predetermined ending condition is fulfilled; I) forming (318) an accepted absolute value set and a sample set by deleting the greatest element from the reference set.

WO 2004/006450 A1